



Mark Scheme (Results)

November 2024

Pearson Edexcel International GCSE  
In Physics (4PH1) Paper 1P

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

November 2024

Question Paper Log Number P75952A

Publications Code 4PH1\_1P\_2411\_MS

All the material in this publication is copyright



© Pearson Education Ltd 2024

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	elliptical orbit drawn; star (approximately) at the focus of the ellipse;	allow partial elliptical orbits reject if star at the centre of the ellipse	2
(b) (i)	arrow drawn pointing from moon towards planet;	arrow does not need to start at moon but line of arrow must pass through moon and planet ignore force arrows associated with comet	1
(ii)	<b>B - gravitational;</b>  A is incorrect as any electrical force would not be strong enough C is incorrect as any magnetic force would not be strong enough D is incorrect as nuclear forces are short range		1
(c)	substitution into given formula;  correct evaluation; answer given to 2 s.f.;	allow use of 3.14, 22/7 for $\pi$ -1 for POT error any final value expressed to 2 s.f. gets this mark	3
	e.g. orbital speed = $2\pi \times 8.7 \times 10^{10} / 2.5 \times 10^8$ orbital speed = 2186.54848... (m/s)  orbital speed = 2200 (m/s)	allow 2185.44, 2187.4... (m/s) allow $2.2 \times 10^3$ (m/s)	
(d)	blue star is hotter (than the Sun);  region is further away (from the star);	allow blue star is hottest allow region for blue star has larger radius / eq	2

Total for Question 1 = 9 marks

Question number	Answer	Notes	Marks
2 (a) (i)	kinetic energy = $\frac{1}{2} \times \text{mass} \times \text{speed}^2$ ;	allow velocity for speed allow standard symbols e.g. $\text{KE} = \frac{1}{2} \times m \times v^2$	1
(ii)	substitution; evaluation;  e.g. $\text{KE} = \frac{1}{2} \times 85 \times 4.5^2$ $\text{KE} = 860 \text{ (J)}$	allow 861, 860.6, 860.62, 860.63, 860.625 (J)	2
(iii)	identical answer to (ii);	expect 860 (J) allow ecf from (ii)	1
(iv)	gravitational potential energy = mass $\times$ g $\times$ height;	allow standard symbols e.g. $\text{GPE} = m \times g \times h$ ignore gravity for g	1
(v)	substitution;  rearrangement; evaluation;  e.g. $860 = 85 \times 10 \times h$ $h = 860 \div 850 = 1.0(125) \text{ (m)}$ (h =) 1.0 (m)	allow use of $g = 9.8, 9.81$ allow ecf from (ii) or (iii)  allow use of $v^2 = u^2 + 2as$ for full marks  allow 1 (m) if supported by valid working allow 1.00 - 1.03 (m)	3
(b) (i)	velocity shown to decrease from an initial positive value;  straight line of negative gradient drawn; finishes at 0 (m/s);  e.g. 	allow if velocity initially increases from 0 (m/s) and then decreases reject if velocity remains at maximum value (flat line) judge by eye  OR 	3

(ii)	<p><b>B - gradient;</b></p> <p>A is incorrect as that would give the distance travelled</p> <p>C and D are incorrect as the x and y axes are integral features of the graph, not the data</p>		1
(iii)	<p>any two from:</p> <p>MP1. idea that Moon and Earth have different masses;</p> <p>MP2. gravitational field strength is proportional to mass;</p> <p>MP3. gravitational field strength on the Moon is less;</p>	<p>allow RA</p> <p>allow density for mass</p> <p>condone Earth is heavier/Moon is lighter</p> <p>allow RA</p> <p>condone less gravity on Moon</p>	2

Total for Question 2 = 14 marks

Question number	Answer	Notes	Marks
3 (a)	plastic is an (electrical) insulator;  idea that it prevents an electric shock;	allow idea the plastic doesn't conduct (electricity) allow reference to double insulation ignore references to current transferring to surroundings	2
(b) (i)	power = current $\times$ voltage;	allow standard symbols and rearrangements e.g. $I = P / V$ ignore C, c for current	1
(ii)	substitution OR rearrangement; evaluation;  e.g. $350 = \text{current} \times 230$ OR $\text{current} = \text{power} / \text{voltage}$ (current =) 1.5 (A)	allow 1.52... (A)	2
(c)	idea the circuit breaker is faster (at isolating circuit); idea the circuit breaker is resettable;	allow circuit breaker is more sensitive allow circuit breaker can be reused / used multiple times	2

Total for Question 3 = 7 marks

Question number	Answer	Notes	Marks																																	
4 (a)	becquerel(s)/Bq;	accept with any prefix (e.g. kBq) or recognisable spelling allow bq, BQ	1																																	
(b)	idea that background radiation is always present;  appropriate named source of background radiation mentioned;	allow radiation all around us, in the surroundings/environment, everywhere etc. e.g. from: <ul style="list-style-type: none"><li>the Sun / cosmic rays</li><li>rocks / soil</li><li>radon</li><li>weapons testing</li><li>food</li><li>nuclear disasters</li><li>medical equipment</li></ul> reject CMBR	2																																	
(c) (i)	(average) time; for halving of: <ul style="list-style-type: none"><li>the (radio)activity;</li><li>number of nuclei;</li><li>mass of isotope;</li></ul>	allow how long it takes  allow count rate for activity allow atoms for nuclei ignore mass of substance/sample	2																																	
(ii)	lines drawn on graph indicating an activity and half of that activity; half-life = 19 (minutes);	e.g. line drawn across from 600 and down to 19 accept any answer that rounds to 19 (minutes) for 2 marks	2																																	
(d)	data point taken from graph; constant correctly calculated using data point; constant correctly calculated from a second data point taken from graph;  conclusion consistent with calculated constants;	<table><tr><th>Time</th><th>Activity</th><th>Constant</th></tr><tr><td>0</td><td>1200</td><td>0</td></tr><tr><td>5</td><td>1000</td><td>5000</td></tr><tr><td>10</td><td>840</td><td>8400</td></tr><tr><td>11-11.5</td><td>800</td><td>8800-9200</td></tr><tr><td>15</td><td>700</td><td>10 500</td></tr><tr><td>19</td><td>600</td><td>11 400</td></tr><tr><td>20</td><td>580</td><td>11 600</td></tr><tr><td>25</td><td>480</td><td>12 000</td></tr><tr><td>30</td><td>400</td><td>12 000</td></tr><tr><td>35</td><td>330-340</td><td>11 550-11 900</td></tr></table> dependent on first three marks being awarded (DOP) conclusion must be clearly linked to a comparison of the constants	Time	Activity	Constant	0	1200	0	5	1000	5000	10	840	8400	11-11.5	800	8800-9200	15	700	10 500	19	600	11 400	20	580	11 600	25	480	12 000	30	400	12 000	35	330-340	11 550-11 900	4
Time	Activity	Constant																																		
0	1200	0																																		
5	1000	5000																																		
10	840	8400																																		
11-11.5	800	8800-9200																																		
15	700	10 500																																		
19	600	11 400																																		
20	580	11 600																																		
25	480	12 000																																		
30	400	12 000																																		
35	330-340	11 550-11 900																																		

Total for Question 4 = 11 marks



Question number	Answer	Notes	Marks
5 (a)	at least three horizontal, straight field lines, parallel and equally spaced; at least one arrow on field line from N to S;	judge by eye  reject any contradictory field line directions for this mark	2
(b)	vertical arrow;  downwards;	allow arrow drawn anywhere on diagram DOP	2
(c)	<b>any three from:</b> MP1. wire/coil/current has a magnetic field; MP2. field of wire/coil/current interacts with field of magnet;  MP3. idea of there being a force <b>on wire AB or CD</b> ;  MP4. idea that <b>forces</b> are in opposite/different directions;  MP5. (because) the currents (in the sides of the coil) are in opposite/different directions;  <b>with one from:</b> MP6. forces do not change direction;  MP7. idea that current cannot swap direction;  MP8. lack of split-ring / commutator;	allow idea that fields overlap ignore cutting field allow if shown on diagram allow upward <b>force</b> on one side and downward <b>force</b> on other side ignore references to movement  allow idea that there is no (net) moment (when the coil is vertical) allow current is not a.c. allow idea that a split-ring / commutator is needed	4

Total for Question 5 = 8 marks

Question number	Answer	Notes	Marks
6 (a)	<p><b>any three from:</b></p> <p>MP1. means of recording position of block;</p> <p>MP2. means of recording rays;</p> <p>MP3. mark normal where light enters/leaves the block;</p> <p>MP4. measurement of angles of incidence and refraction;</p> <p>MP5. repeat for range of angles of incidence;</p> <p><b>with any two from:</b></p> <p>MP6. <math>\sin(i) \div \sin(r)</math> used to calculate <math>n</math>;</p> <p>MP7. for different angles of incidence and averaged;</p>	<p>allow drawing around block on paper</p> <p>allow drawing paths of rays, marking positions of incident/emerging rays</p> <p>allow if shown in diagram</p> <p>allow graph plotted of <math>\sin(i)</math> against <math>\sin(r)</math></p> <p>allow use of gradient to determine <math>n</math></p>	5
(b) (i)	$\sin(c) = 1/n$ ;	allow any correct rearrangement allow word formula	1
(ii)	<p>substitution OR rearrangement; evaluation;</p> <p>e.g.  <math>\sin(c) = 1 / 1.2</math> OR <math>c = \sin^{-1}(1/n)</math>  <math>(c = \sin^{-1}(1/1.2) =) 56</math> (degrees)</p>	<p>allow <math>\sin(c) = 0.83...</math></p> <p>allow 56.4... (degrees)</p> <p>allow answers that round to 56 (degrees)</p>	2
(iii)	<p>any two from:</p> <p>MP1. idea that all light is reflected inside a medium;</p> <p>MP2. for light travelling from a higher refractive index to a lower refractive index;</p> <p>MP3. (when) angle (of incidence) is greater than critical angle;</p>	allow if seen in a diagram allow idea that all light is reflected rather than being refracted allow higher (optical) density to lower (optical) density	2
(c) (i)	24 (degrees);	allow 22-26 (degrees)	1
(ii)	<p>ray emerges into air;</p> <p>ray refracts correctly, bending away from the normal;</p>	<p>allow dashed lines</p> <p>arrow not required</p> <p>judge by eye</p> <p>ignore reflected rays</p>	2

Total for Question 6 = 13 marks


Question number	Answer	Notes	Marks
7	(a) (i)	weight / gravitational (force) / gravitational pull (of Earth);	1
	(ii)	one vertical upwards arrow and one vertical downwards arrow;  two arrows are equal length;	2
(b)	(i)	pressure (difference) = density $\times$ g $\times$ height;	1
	(ii)	substitution;  evaluation in kPa;  e.g. pressure = $1030 \times 10 \times 38$ (pressure =) 390 (kPa)	2
(c)	(i)	idea of collisions between particles and <b>the walls</b> of the storage tank;  idea of particles exerting a force on the walls;  idea of a force exerted over an area gives a pressure;	3

(ii)	<p>conversion of either temperature in kelvin; substitution into given formula;</p> <p>rearrangement; evaluation;</p> <p>e.g.  <math>T = 275.5 \text{ (K)}</math> or <math>T = 291 \text{ (K)}</math>  <math>410 / 275.5 = p_2 / 291</math>  <math>p_2 = 410 \times 291 / 275.5</math>  <math>(p_2 =) 430 \text{ (kPa)}</math></p>	<p>allow if seen anywhere in working allow ecf for °C values used or incorrect kelvin temperatures accept calculation of constant value i.e. <math>410 \div 275.5 = 1.488...</math></p> <p>–1 for POT error</p> <p>3000, 2952 (kPa) scores 3 marks (not converting to kelvin)</p> <p>allow 433.1, 433.0... (kPa)</p>	4
------	--	--	---

Total for Question 7 = 13 marks


Question number	Answer	Notes	Marks
8 (a)	reference to vibrations / oscillations;  (vibrations/oscillations) are parallel to direction of wave/energy transfer for longitudinal AND (vibrations/oscillations) are perpendicular to direction of wave/energy transfer for transverse;	both marking points can be awarded from clearly labelled diagrams ignore up and down, forwards and backwards allow equivalent statements of parallel/perpendicular	2
(b)	wavefronts (at whale B) are compressed / wavelength (at whale B) is smaller; speed of sound is constant; frequency = speed of sound ÷ wavelength (therefore frequency increases);	allow alternative phrases for compressed  allow any rearrangement of the formula allow symbols for formula	3
(c)	substitution into speed = frequency × wavelength; rearrangement; evaluation;  e.g. $3.0 \times 10^8 = \text{frequency} \times 5.4 \times 10^{-7}$ $\text{frequency} = 3.0 \times 10^8 / 5.4 \times 10^{-7}$ (frequency =) $5.6 \times 10^{14}$ (Hz)	-1 for POT error if no other marks scored, allow selection of correct formula for 1 mark  allow $6 \times 10^{14}$ , $5.55... \times 10^{14}$ (Hz) reject $5.5 \times 10^{14}$ (Hz)	3

Total for Question 8 = 8 marks

Question number	Answer	Notes	Marks
9 (a)	<p>complete circuit with correct symbols for cell, ammeter and voltmeter;</p> <p>variable resistor in series with diode;</p> <p>ammeter in series with diode;</p> <p>voltmeter in parallel with diode;</p> <p>e.g.</p> 	<p>condone more than one cell / battery drawn</p> <p>ignore orientation of diode relative to cell</p> <p>circuit symbol must be correct</p> <p>condone incorrect orientation of symbol</p> <p>reject if voltmeter drawn in parallel with variable resistor and diode</p>	4
(b) (i)	0.38 (A);		1
(ii)	voltage = current $\times$ resistance;	<p>allow standard symbols and rearrangements</p> <p>e.g. <math>R = V / I</math></p> <p>ignore c, C for current</p>	1
(iii)	<p>substitution OR rearrangement; evaluation;</p> <p>e.g.</p> <p><math>2.9 = 0.15 \times R</math> OR <math>R = V / I</math></p> <p>(resistance =) 19 (<math>\Omega</math>)</p>	<p>allow 19.3... (<math>\Omega</math>)</p>	2
(iv)	<p>MP1. calculation of resistance in diagram 2;</p> <p>MP2. idea that resistance is larger in diagram 2;</p> <p>MP3. calculation of current in diagram 2;</p>	<p>allow <math>19(.3) + 13</math> or <math>32(.3)</math> (<math>\Omega</math>) seen</p> <p>allow ecf from (iii)</p> <p>allow <math>12.6 \Omega</math> used for <math>R_B</math></p> <p>allow <math>R_A</math></p> <p>award this mark if MP1 awarded</p> <p>allow ecf from MP1</p>	4

	MP4. conclusion that current is smaller in diagram 2;	e.g. $I (= 2.9 / 32(.3)) = 0.09 \text{ (A)}$ dependent on one of MP1-MP3 being awarded allow RA	
--	---	---	--

Total for Question 9 = 12 marks

Question number	Answer	Notes	Marks										
10 (a)	(i) metal/material;	ignore unqualified “type of rod”	1										
	(ii) time (taken for coin to fall off);	allow how long it takes (for the coin to fall off)	1										
	(iii) any two from: MP1. rods the same length; MP2. rods the same diameter/thickness/width/area; MP3. coins of the same mass; MP4. same mass/amount of wax;	allow a mark for “rods same size/volume” if MP1 or MP2 have not been awarded  allow same type of wax	2										
	(iv) repeat the experiment;  calculate the mean (time for each rod);	reject if method modified e.g. testing additional metals allow calculate the average allow identify/ignore anomalies	2										
	(b) (i)	suitable linear scale for time axis;  time axis labelled with quantity and unit AND material axis labelled;  all bars correct length to within ½ small square;  e.g. 	reject multiples of 30, 45, 60, 80 for 10 small squares allow use of false origin scale should be such that length of tallest bar must be at least half the grid condone missing “material” label if bars are labelled iron, steel etc. <table><tr><th>Material</th><th>Time for coin to drop in seconds</th></tr><tr><td>iron</td><td>180</td></tr><tr><td>steel</td><td>230</td></tr><tr><td>aluminium</td><td>140</td></tr><tr><td>copper</td><td>90</td></tr></table>	Material	Time for coin to drop in seconds	iron	180	steel	230	aluminium	140	copper	90
Material	Time for coin to drop in seconds												
iron	180												
steel	230												
aluminium	140												
copper	90												
	(ii) material/data is categoric/discrete/not continuous;	allow discontinuous for not continuous	1										

Total for Question 10 = 10 marks



Question number	Answer	Notes	Marks												
11 (a)	<p><b>D - nuclear fusion;</b></p> <p>A is incorrect as the abundance of nuclei with high proton number is too small B is incorrect as the energy release per decay is too small C is incorrect as fusion below Fe-57 is endothermic</p>		1												
(b)	<p>all numbers written in the correct rows = 4 marks;;;; any three numbers written in the correct rows = 3 marks;;;; any two numbers written in the correct rows = 2 marks;; any one number written in the correct row = 1 mark; e.g.</p> <table><tr><th>Stages of development</th><th>Order number</th></tr><tr><td>a black hole is formed</td><td>5</td></tr><tr><td>heavy elements are made in the core of the star</td><td>3</td></tr><tr><td>the star becomes a red supergiant</td><td>2</td></tr><tr><td>the star explodes in a supernova</td><td>4</td></tr><tr><td>the core of the star runs out of hydrogen</td><td>1</td></tr></table>	Stages of development	Order number	a black hole is formed	5	heavy elements are made in the core of the star	3	the star becomes a red supergiant	2	the star explodes in a supernova	4	the core of the star runs out of hydrogen	1		4
Stages of development	Order number														
a black hole is formed	5														
heavy elements are made in the core of the star	3														
the star becomes a red supergiant	2														
the star explodes in a supernova	4														
the core of the star runs out of hydrogen	1														

Total for Question 11 = 5 marks

